

Seasonal forecasts

presented by:

Willem.Landman@up.ac.za
WALandman1981@gmail.com



Seasonal Forecast Worx



tinyurl.com/SeasonalForecastWorx

UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA

Seasonal Climate Forecasts

Latest Update: 15 August 2025

- The seasonal forecasts presented here by **Seasonal Forecast Worx** are based on forecast output of the coupled ocean-atmosphere models administered through the North American Multi-Model Ensemble (NMME) prediction experiment (<http://www.cpc.ncep.noaa.gov/products/NMME/>; Kirtman et al. 2014). NMME real-time seasonal forecast and hindcast (re-forecast) data are obtained from the data library (<http://iridl.ldeo.columbia.edu/>) of the International Research Institute for Climate and Society (IRI; <http://iri.columbia.edu/>).
- NMME forecasts are routinely produced and are statistically improved and tailored for southern Africa and for global sea-surface temperatures by employees and post-graduate students in the Department of Geography, Geoinformatics and Meteorology at the University of Pretoria (<http://www.up.ac.za/en/geography-geoinformatics-and-meteorology/>). Statistical post-processing is performed with the CPT software (<http://iri.columbia.edu/our-expertise/climate/tools/cpt/>).
- Why do we apply statistical methods to climate model forecasts?
- “...**statistical correction methods treating individual locations (e.g. multiple regression or principal component regression) may be recommended for today’s coupled climate model forecasts**”. (Barnston and Tippett, 2017).
- Why do we not use just a single model in our forecasts?
- “...**multi-model forecasts outperform the single model forecasts...**” (Landman and Beraki, 2012).
- For the official seasonal forecast for South Africa, visit the South African Weather Service website at: <https://www.weathersa.co.za/home/seasonalclimate>

ENSO and Global SST Forecasts


Skilful seasonal forecasts are becoming increasingly vital tools for
climate change adaptation in southern Africa

Prediction Method

- Forecasts for global sea-surface temperature (SST) fields are obtained through a combination of NMME models and a linear statistical model, that uses antecedent SST as a predictor (Landman et al. 2011). Forecasts for the Niño3.4 area (see insert) are derived from the global forecasts.
- SST forecasts from the NMME models are variance and bias corrected.



El Niño and Rainfall

El Niño conditions in the tropical Pacific are known to shift rainfall patterns in many different parts of the world. Although they vary somewhat from one El Niño to the next, the strongest shifts remain fairly consistent in the regions and seasons shown on the map below.




La Niña and Rainfall

La Niña conditions in the tropical Pacific are known to shift rainfall patterns in many different parts of the world. Although they vary somewhat from one La Niña to the next, the strongest shifts remain fairly consistent in the regions and seasons shown on the map below.



CSiriMM Nino3.4 SST Forecast

Issued on: 14-Aug-2025




Middle red line: the forecast
Thin red lines: 25% confidence levels

Verification of these forecasts: <https://tinyurl.com/UPENSOforecastVerification>

Oceanic Niño Index (ONI)


http://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml




IC: the month in which the forecast was made

SST anomalies (in °C, where blue is cooler and orange is warmer)


SST ASO IC:Aug2025




SST SON IC:Aug2025




SST OND IC:Aug2025




SST NDJ IC:Aug2025



SST DJF IC:Aug2025





SST JFM IC:Aug2025



Round-up: ENSO

- The UP model (on previous pages) is predicting mostly ENSO-neutral conditions throughout the austral summer. However, some of the international models are predicting a weak La Niña event



The UP model forms part of this plume, and is marked as "CS-IRI-MM"

Africa forecasts, south of 20°N

Prediction Method

- Three-month seasons for seasonal rainfall totals and average maximum temperatures of NMME ensemble mean forecasts are recalibrated to the Climatic Research Unit (CRU; Harris et al. 2014) grids ($0.5^{\circ} \times 0.5^{\circ}$). Probabilistic forecasts are subsequently produced from the error variance obtained from a 5-year-out cross-validation process (Troccoli et al. 2008). Forecasts cover a 6-month period.
- Forecasts are produced for three categories:
 - **Above:** Above-normal (“wet” rainfall totals / “hot” maximum temperatures higher than the 75th percentile of the climatological record)
 - **Below:** Below-normal (“dry” rainfall totals / “cool” maximum temperatures lower than the 25th percentile of the climatological record)
 - **Normal:** Near-normal (“average” season)
- Verification of forecast performance:
 - ROC Area (Below-Normal) – The forecast system’s ability to discriminate dry or cool seasons from the rest of the seasons over a 23-year test period. ROC values should be higher than 0.5 for a forecast system to be considered skilful.
 - ROC Area (Above-Normal) – The forecast system’s ability to discriminate wet or hot seasons from the rest of the seasons over a 23-year test period. ROC values should be higher than 0.5 for a forecast system to be considered skilful.
 - The white areas on the forecast maps
 - No forecast - forecasts for the near-normal category do not have skill and are therefore not shown

Forecasts are probabilistic


Probabilistic forecasts can help users understand risks and opportunities (forewarned is forearmed) in order to make more informed decisions.

The seasonal rainfall and maximum temperature forecasts to follow are expressed in probabilities, shown as the % chance of the most likely outcome of 3 categories. The colour of the scale reflects the most likely category and the % shows the probability of that outcome. Only ONE of the ROC area skill assessment maps should be consulted, depending on the category shown on the forecast map (Above- or Below-Normal), and the higher the ROC value, the more skilful the forecast for that pixel is. The probabilities shown are always less than 100% - so there is no absolute certainty that the less favoured outcome will not occur. For example, if the forecast claims a 75% chance of below-normal rainfall totals for a season (i.e. drought), it means that 1 out of 4 times it will **not** develop into a drought.

The nature of a probabilistic forecast implies that the less likely outcomes are always possible. In fact, for the probabilistic forecasts to be considered reliable, the less likely outcomes will and must occasionally occur.


Note: Probabilistic forecasts are considered reliable when the forecast probability is an accurate estimation of the relative frequency of the predicted outcome. In other words, forecasts are reliable if the observation falls within the category (Below-, Near- or Above-Normal) as frequently as the forecast implies

ASO 2025 Rainfall; ICs: Aug




Legend

Rainfall Prob




No forecast

ROC Area (Above-Normal): ASO Rainfall




Legend

ROC




ROC Area (Below-Normal): ASO Rainfall




Legend

ROC




SON 2025 Rainfall; ICs: Aug




Legend

Rainfall Prob




□ No forecast

ROC Area (Above-Normal): SON Rainfall




Legend

ROC




ROC Area (Below-Normal): SON Rainfall




Legend

ROC




OND 2025 Rainfall; ICs: Aug




Legend

Rainfall Prob



□ No forecast

ROC Area (Above-Normal): OND Rainfall




Legend

ROC

0.8 - 0.9
0.7 - 0.8
0.6 - 0.7
0.5 - 0.6
0 - 0.5

ROC Area (Below-Normal): OND Rainfall




Legend

ROC


0.8 - 0.9
0.7 - 0.8
0.6 - 0.7
0.5 - 0.6
0 - 0.5

NDJ 2025/26 Rainfall; ICs: Aug




Legend

Rainfall Prob




□ No forecast

ROC Area (Above-Normal): NDJ Rainfall



ROC Area (Below-Normal): NDJ Rainfall




Round-up: Rainfall south of 15°S


- Spring rainfall over the SW Cape and parts of Namibia is likely to be above-normal, but below-normal over the eastern and northern parts
- In general, chances of above-normal rainfall increase in the summer rainfall region towards early and mid-summer

Observed Rainfall

Rainfall (% of normal): July 2025
July long-term mean: 1981–2010



Rainfall (% of normal): May–June–July 2025
May–June–July long-term mean: 1981–2010




Recorded rainfall for July and the May–June–July season show below-normal rainfall over the brown areas and above-normal rainfall over the green areas

Maps prepared by Dr. Christien Engelbrecht


Compare these OLD forecasts with the observations on the previous page




MJJ 2025 Rainfall; ICs: Feb




MJJ 2025 Rainfall; ICs: Mar




MJJ 2025 Rainfall; ICs: Apr



MJJ 2025 Rainfall; ICs: May




ASO 2025 Max Temp; ICs: Aug




Legend

Max Temp Prob




No Forecast

ROC Area (Above-Normal): ASO Max Temp




Legend

ROC




ROC Area (Below-Normal): ASO Max Temp




Legend


ROC




SON 2025 Max Temp; ICs: Aug




ROC Area (Above-Normal): SON Max Temp




ROC Area (Below-Normal): SON Max Temp



OND 2025 Max Temp; ICs: Aug




Legend




No Forecast


ROC Area (Above-Normal): OND Max Temp




Legend




ROC Area (Below-Normal): OND Max Temp




Legend




NDJ 2025/26 Max Temp; ICs: Aug



ROC Area (Above-Normal): NDJ Max Temp



ROC Area (Below-Normal): NDJ Max Temp



Round-up: South of 15°S Max Temp


- Below-normal maximum temperatures are expected in late winter over SW Cape
- Above-normal maximum temperatures are predicted over the larger part of the region at first, but below-normal maximum temperatures are expected towards mid-summer in anticipation of an expected wet summer, except over the southern coastal areas where it is more likely to be above-normal

References

- Barnston, A.G. and Tippett, M.K., 2017: Do statistical pattern corrections improve seasonal climate predictions in the North American Multimodel Ensemble models? *Journal of Climate*, 30: 8335-8355. doi: 10.1175/JCLI-D-17-0054.1
- Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H., 2014: Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology*, 34: 623-642. doi: 10.1002/joc.3711
- Kirtman, B. P. and Co-authors 2014: The North American Multimodel Ensemble: Phase-1 seasonal-to-interannual prediction; Phase-2 toward developing intraseasonal prediction. *Bulletin of the American Meteorological Society*. 95, 585–601. doi: <http://dx.doi.org/10.1175/BAMS-D-12-00050.1>
- Landman, W.A., and Beraki, A., 2012: Multi-model forecast skill for midsummer rainfall over southern Africa. *International Journal of Climatology*, 32: 303-314. doi: 10.1002/joc.2273.
- Landman, W.A., Archer, E. and Tadross, M., 2016: Decision-relevant information on seasonal time scales – the case of a farm in northern Namibia. *Conference Proceedings of the 32nd Annual Conference of the South African Society for Atmospheric Science*, Cape Town, 31 October to 1 November 2016, pp 69-72. ISBN 978-0-620-72974-1.
- Landman, W.A., Archer, E. and Tadross, M. (2019): How costly are poor seasonal forecasts? Peer reviewed abstracts, 35th Annual conference of the South African Society for Atmospheric Science, Vanderbijlpark, 8 to 9 October 2019, pp 60-63. ISBN 978-0-6398442-0-6.
- Landman, W.A., Archer, E.R.M and Tadross, M.A (2020a). Citizen science for the prediction of climate extremes in South Africa and Namibia. *Frontiers in Climate*, 2:5, doi: 10.3389/fclim.2020.00005
- Landman, W.A., DeWitt, D., and Lee, D.-E., 2011: The high-resolution global SST forecast set of the CSIR. *Conference Proceedings of the 27th Annual Conference of South African Society for Atmospheric Sciences*, 22-23 September 2011, Hartbeespoort, North-West Province, South Africa. ISBN 978-0-620-50849-0
- Landman, W.A., Sweijd, N., Masedi, N. Minakawa, N. (2020b). The development and prudent application of climate-based forecasts of seasonal malaria in the Limpopo province in South Africa. *Environmental Development*, 35, 100522, doi: 10.1016/j.envdev.2020.100522.
- Landman, W.A., Tadross, M., Archer, E., Johnston, P. (2023). Probabilistic vs deterministic forecasts - interpreting skill statistics for the benefit of users. *Water SA*, 49(3), 192–198. <https://doi.org/10.17159/wsa/2023.v49.i3.4058>
- Landman, W.A., DeWitt, D. Lee, D.-E., Beraki, A. and Lötter, D., 2012: Seasonal rainfall prediction skill over South Africa: 1- vs. 2-tiered forecasting systems. *Weather and Forecasting*, 27: 489-501. DOI: 10.1175/WAF-D-11-00078.1
- Muchuru, S., Landman, W.A. and DeWitt, D., 2016: Prediction of inflows into Lake Kariba using a combination of physical and empirical models. *International Journal of Climatology*, 36: 2570–2581, DOI: 10.1002/joc.4513.
- Troccoli, A., Harrison, M., Anderson, D.L.T. and Mason, S.J., 2008: *Seasonal Climate: Forecasting and Managing Risk*. NATO Science Series on Earth and Environmental Sciences, Vol. 82, Springer, 467 pp.

Financial support from:

- Wellcome Trust “Climatefocused Analytics and Modelling for Mosquito-borne Infections in Southern Africa” (CAMMISA; 2025 to 2029)
- The National Research Foundation
 - Incentive Funding for Rated Researchers (since 2017)
 - Project: Application of knowledge for the management of extreme climate events (APECX; 2022 to 2025)
- ACCESS (Alliance for Collaboration on Climate and Earth System Science) through the project “Investigating predictability of seasonal anomalies for societal benefit” (2018 to 2021)
- Water Research Commission through administering the international project “Research-based Assessment of Integrated approaches to Nature-based SOLUTIONS (RainSolutions)” (2020 to 2022)



The forecast is produced by Prof Willem Landman of the University of Pretoria, South Africa, and issued on or around the 15th of each month. Please feel free to contact me at WALandman1981@gmail.com

Acknowledgments to Dr Peter Johnston of the University of Cape Town for professional comments and advice

Disclaimer: The author has compiled this forecast guidance as a service to users for application in appropriate sectors, but cannot be held responsible for inaccuracies contained therein

Student participation in forecast system development



Stephanie Hinze, BSc (Honours)(Meteorology):

Statistical downscaling using large and high-resolution data sets, forecast displays for SADC rainfall and maximum temperatures, forecast verification



Surprise Mhlongo, BSc (Honours)(Meteorology):

Improving on SST forecast system through pattern correction, correlation vs covariance approaches, forecast output combination (multi-model approaches), mean and bias correction, and correct for skill



Shepherd Muchuru, PhD (Meteorology):

Statistical modelling to relate large-scale features to seasonal inflows into Lake Kariba in southern Africa. Two predictions systems: 1) using antecedent seasonal rainfall totals over the upper Zambezi catchment as predictor in a baseline model, and 2) using predicted low-level atmospheric circulation of a coupled ocean–atmosphere general circulation model as predictor.



Pearl Gosiame, BSc (Honours)(Meteorology):

Development of hydro-climate predictions models for dam levels and downstream flows of the Vaal Dam. Predictors considered include historical rainfall over the catchment, SST and output from global climate models.



Idani Mandiwana, BSc (Honours) (Meteorology):

Seasonal rainfall forecast verification of real-time forecasts produced by SFW over the 5-year period from 2018 to 2022. Area is SADC south of 17° South.



Aimee Serafini, MSc (Meteorology): Development of seasonal forecast systems for farmers in Namibia using farm observations and output from global climate models.