

Seasonal forecasts

presented by:

Tel: +27(82)644-5304
Willem.Landman@up.ac.za



<https://tinyurl.com/ybrb3a72>



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA






Seasonal Forecast Worx

Latest Update: 10 November 2020

Are you a farmer who wants to make use of science-based seasonal predictions for your farm? If you are interested to be part of an initiative at the University of Pretoria that involves the development of seasonal forecast systems for farms, specifically tailored to farmers' needs, please send an email to WALandman1981@gmail.com

Have a look at this example of end-of-season crop yield forecasts for a farm near Bapsfontein. The farmer provided several decades of crop-yield data and these data were subsequently used to create a crop forecast model specific to the farm



Above is the crop-yield forecast for the coming season. The forecast is for enhanced probabilities of **below-normal** (low) crop yield for the farm. The farmer may be able with support to use this forecast information to plan for the coming season

On the left are time series of forecast and observed crop yields at the time of harvest for the years indicated. Next to the time series are probabilistic forecasts over the same 9-years for below- (low yields), near- (about average) and above-normal (high yields). For example, in 2008 the forecast and observed index values are high and positive (figure on the left), and the highest predicted probability is for above-normal yield (figure in the middle).


Share your data and become part of this initiative

- The seasonal forecasts presented here by **Seasonal Forecast Worx** are based on forecast output of the coupled ocean-atmosphere models administered through the North American Multi-Model Ensemble (NMME) prediction experiment (<http://www.cpc.ncep.noaa.gov/products/NMME/>; Kirtman et al. 2014). NMME real-time seasonal forecast and hindcast (re-forecast) data are obtained from the data library (<http://iridl.ldeo.columbia.edu/>) of the International Research Institute for Climate and Society (IRI; <http://iri.columbia.edu/>).
- NMME forecasts are routinely produced and are statistically improved and tailored for southern Africa and for global sea-surface temperatures by employees and post-graduate students in the Department of Geography, Geoinformatics and Meteorology at the University of Pretoria (<http://www.up.ac.za/en/geography-geoinformatics-and-meteorology/>). Statistical post-processing is performed with the CPT software (<http://iri.columbia.edu/our-expertise/climate/tools/cpt/>).
- Why do we apply statistical methods to climate model forecasts?
 “...**statistical correction methods treating individual locations (e.g. multiple regression or principal component regression) may be recommended for today’s coupled climate model forecasts**”. (Barnston and Tippett, 2017).
- Why do we not use just a single model in our forecasts?
 “...**multi-model forecasts outperform the single model forecasts...**” (Landman and Beraki, 2012).
- For the official seasonal forecast for South Africa, visit the South African Weather Service website at <http://www.weathersa.co.za/images/data/longrange/gfcsa/scw.pdf>

ENSO and Global SST Forecasts


Prediction Method

- Forecasts for global sea-surface temperature (SST) fields are obtained through a combination of NMME models and a linear statistical model, that uses antecedent SST as a predictor (Landman et al. 2011). Forecasts for the Niño3.4 area (see insert) are derived from the global forecasts.
- SST forecasts from the NMME models are variance and bias corrected.
- Three-month Niño3.4 SST forecasts are produced for three categories:
 - **El Niño:** SST above the 75th percentile
 - **La Niña:** SST below the 25th percentile
 - **Neutral:** Neither El Niño nor La Niña




CSiriMM Nino3.4 SST Forecast

Issued on: 10-Nov-2020




Oceanic Niño Index (ONI)

http://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml




SST anomalies


SST NDJ IC:Nov2020




SST DJF IC:Nov2020




SST JFM IC:Nov2020




SST FMA IC:Nov2020



SST MAM IC:Nov2020




SST AMJ IC:Nov2020



Round-up: ENSO

- The UP model predicts weak to moderate La Niña conditions for mid-summer, but ENSO-neutral conditions towards autumn
- Most forecast models continue to be for colder SST, with just a few warmer than the UP model (CS-IRI-MM)



Southern Africa Forecasts

Prediction Method

- Three-month seasons for seasonal rainfall totals and average maximum temperatures of NMME ensemble mean forecasts are interpolated to Climatic Research Unit (CRU; Harris et al. 2014) grids ($0.5^{\circ} \times 0.5^{\circ}$), by correcting the mean and variance biases of the NMME forecasts. Probabilistic forecasts are subsequently produced from the error variance obtained from a 5-year-out cross-validation process (Troccoli et al. 2008). Forecasts cover a 6-month period.
- Forecasts are produced for three categories:
 - **Above:** Above-normal (“wet” / “hot”, rainfall totals / maximum temperatures higher than the 75th percentile of the climatological record)
 - **Below:** Below-normal (“dry” / “cool”, rainfall totals / maximum temperatures lower than the 25th percentile of the climatological record)
 - **Normal:** Near-normal (“average” season)
- Verification:
 - ROC Area (Below-Normal) – The forecast system’s ability to discriminate dry or cool seasons from the rest of the seasons over a 32-year test period. ROC values should be higher than 0.5 for a forecast system to be considered skilful.
 - ROC Area (Above-Normal) – The forecast system’s ability to discriminate wet or hot seasons from the rest of the seasons over a 32-year test period. ROC values should be higher than 0.5 for a forecast system to be considered skilful.

Forecasts are probabilistic


Probabilistic forecasts can help users understand risks and opportunities (forewarned is forearmed) in order to make more informed decisions.

The seasonal rainfall and maximum temperature forecast to follow are probabilities (% chance) of only the most likely outcome for below-, near-, or above-normal (B, N or A). The probabilities shown are always less than 100% - so there is no absolute certainty that the less favoured outcome will not occur. For example, if the forecast claims a 75% of below-normal rainfall totals for a season (i.e. drought), it means that 1 out of 4 times it will not develop into a drought.


The nature of a probabilistic forecast implies that the less likely outcomes are always possible. In fact, for the probabilistic forecasts to be considered reliable, the less likely outcomes will and must occasionally occur.

Note: Probabilistic forecasts are considered reliable when the forecast probability is an accurate estimation of the relative frequency of the predicted outcome. In other words, forecasts are reliable if the observation falls within the category (B, N or A) as frequently as the forecast implies


NDJ 2020/21 Rainfall; ICs: Nov




ROC Area (Above-Normal): NDJ Rainfall




ROC Area (Below-Normal): NDJ Rainfall




DJF 2020/21 Rainfall; ICs: Nov




ROC Area (Above-Normal): DJF Rainfall




ROC Area (Below-Normal): DJF Rainfall




JFM 2021 Rainfall; ICs: Nov




ROC Area (Above-Normal): JFM Rainfall



ROC Area (Below-Normal): JFM Rainfall




FMA 2021 Rainfall; ICs: Nov




Legend

----- Catchment Area
— Main Rivers


Rainfall Prob



ROC Area (Above-Normal): FMA Rainfall



ROC Area (Below-Normal): FMA Rainfall




Round-up: SADC Rainfall

- Favourable rainfall outcomes are expected over the larger part of the forecast region during spring and summer, except over some of the northern parts

Observed SADC Rainfall


Rainfall (% of normal): August–September–October 2020

Relative to August–September–October 1981–2010 rainfall





Rainfall (% of normal): October 2020

October long-term mean: 1981–2010




Maps prepared by Dr. Christien Engelbrecht


NDJ 2020/21 Max Temp; ICs: Nov




ROC Area (Above-Normal): NDJ Max Temp




ROC Area (Below-Normal): NDJ Max Temp




DJF 2020/21 Max Temp; ICs: Nov




ROC Area (Above-Normal): DJF Max Temp




ROC Area (Below-Normal): DJF Max Temp




JFM 2021 Max Temp; ICs: Nov




ROC Area (Above-Normal): JFM Max Temp




ROC Area (Below-Normal): JFM Max Temp




FMA 2021 Max Temp; ICs: Nov



ROC Area (Above-Normal): FMA Max Temp



ROC Area (Below-Normal): FMA Max Temp



Round-up: SADC Max Temp


- Cooler maximum temperatures are likely in association with the increased likelihood of a wet summer season over parts of the region

Tailored Forecasts


1. Bapsfontein end-of-season-yield three-category probabilistic forecast for 2021
2. Probabilistic three-category rainfall forecast for the farm of Robbie Kingsley for Dec-Jan-Feb 2020/21
3. Probabilistic rainfall forecast for Jan-Feb-Mar 2021 for the farm Buschbrunnen near Grootfontein, Namibia
4. Probabilistic three-category malaria forecast for Limpopo for Dec-Jan-Feb 2020/21
5. Probability of exceedance Dec-Jan-Feb 2020/21 inflow forecast for Lake Kariba, Zambia/Zimbabwe
6. Probability of exceedance Jan-Feb-Mar, Feb-Mar-Apr and Mar-Apr-May 2021 downstream flow forecasts for Vaal Dam

Crop-yield data and forecasts for a farm near Bapsfontein, South Africa

Landman et al. (2019)







Re-forecasts for end-of-season crop yields




Dec-Jan-Feb 2020/21 rainfall forecast for farm in the Smithfield district (see map). Rainfall data provided by the farmer, Mr. Robbie Kingsley

Landman et al. (2020a)



JFM rainfall forecast for the farm Buschbrunnen near Grootfontein, Namibia


Landman et al. (2016)




Malaria forecast

Landman et al. (2020b)

Hindcasts



DJF Malaria Forecasts for Limpopo (Issued in Nov)






Retrospective

REAL-TIME

Inflow forecast for Lake Kariba: onset season of DJF

Muchuru et al. (2016)



Current Project, administered by the WRC: *RainSolutions*

Vaal Dam Downstream Flow Predictions







Figure 1
Location of Vaal Dam in Upper Vaal Water Management Area in South Africa (Modified from DWA, 2010)




References

- Barnston, A.G. and Tippett, M.K., 2017: Do statistical pattern corrections improve seasonal climate predictions in the North American Multimodel Ensemble models? *Journal of Climate*, 30: 8335-8355. doi: 10.1175/JCLI-D-17-0054.1
- Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H., 2014: Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology*, 34: 623-642. doi: 10.1002/joc.3711
- Kirtman, B. P. and Co-authors 2014: The North American Multimodel Ensemble: Phase-1 seasonal-to-interannual prediction; Phase-2 toward developing intraseasonal prediction. *Bulletin of the American Meteorological Society*. 95, 585–601. doi: <http://dx.doi.org/10.1175/BAMS-D-12-00050.1>
- Landman, W.A., and Beraki, A., 2012: Multi-model forecast skill for midsummer rainfall over southern Africa. *International Journal of Climatology*, 32: 303-314. doi: 10.1002/joc.2273.
- Landman, W.A., Archer, E. and Tadross, M., 2016: Decision-relevant information on seasonal time scales – the case of a farm in northern Namibia. *Conference Proceedings of the 32nd Annual Conference of the South African Society for Atmospheric Science*, Cape Town, 31 October to 1 November 2016, pp 69-72. ISBN 978-0-620-72974-1.
- Landman, W.A., Archer, E. and Tadross, M. (2019): How costly are poor seasonal forecasts? Peer reviewed abstracts, 35th Annual conference of the South African Society for Atmospheric Science, Vanderbijlpark, 8 to 9 October 2019, pp 60-63. ISBN 978-0-6398442-0-6.
- Landman, W.A., Archer, E.R.M and Tadross, M.A (2020a). Citizen science for the prediction of climate extremes in South Africa and Namibia. *Frontiers in Climate*, 2:5, doi: 10.3389/fclim.2020.00005
- Landman, W.A., DeWitt, D., and Lee, D.-E., 2011: The high-resolution global SST forecast set of the CSIR. *Conference Proceedings of the 27th Annual Conference of South African Society for Atmospheric Sciences*, 22-23 September 2011, Hartbeespoort, North-West Province, South Africa. ISBN 978-0-620-50849-0
- Landman, W.A., Sweijd, N., Masedi, N. Minakawa, N. (2020b). The development and prudent application of climate-based forecasts of seasonal malaria in the Limpopo province in South Africa. *Environmental Development*, 35, 100522, doi: 10.1016/j.envdev.2020.100522.
- Landman, W.A., DeWitt, D. Lee, D.-E., Beraki, A. and Lötter, D., 2012: Seasonal rainfall prediction skill over South Africa: 1- vs. 2-tiered forecasting systems. *Weather and Forecasting*, 27: 489-501. DOI: 10.1175/WAF-D-11-00078.1
- Muchuru, S., Landman, W.A. and DeWitt, D., 2016: Prediction of inflows into Lake Kariba using a combination of physical and empirical models. *International Journal of Climatology*, 36: 2570–2581, DOI: 10.1002/joc.4513.
- Troccoli, A., Harrison, M., Anderson, D.L.T. and Mason, S.J., 2008: *Seasonal Climate: Forecasting and Managing Risk*. NATO Science Series on Earth and Environmental Sciences, Vol. 82, Springer, 467 pp.

Financial support from...

- The National Research Foundation through the Incentive Funding for Rated Researchers
- ACCESS (Alliance for Collaboration on Climate and Earth System Science) through the project “Investigating predictability of seasonal anomalies for societal benefit”
- Water Research Commission through administering the international project “Research-based Assessment of Integrated approaches to Nature-based SOLUTIONS (RainSolutions)”



Student participation in forecast system development



Stephanie Hinze, BSc (Honours)(Meteorology):

Statistical downscaling using large and high-resolution data sets, forecast displays for SADC rainfall and maximum temperatures, forecast verification



Surprise Mhlongo, BSc (Honours)(Meteorology):

Improving on SST forecast system through pattern correction, correlation vs covariance approaches, forecast output combination (multi-model approaches), mean and bias correction, and correct for skill



Shepherd Muchuru, PhD (Meteorology):

Statistical modelling to relate large-scale features to seasonal inflows into Lake Kariba in southern Africa. Two predictions systems: 1) using antecedent seasonal rainfall totals over the upper Zambezi catchment as predictor in a baseline model, and 2) using predicted low-level atmospheric circulation of a coupled ocean–atmosphere general circulation model as predictor.



Pearl Gosiame, BSc (Honours)(Meteorology):

Development of hydro-climate predictions models for dam levels and downstream flows of the Vaal Dam. Predictors considered include historical rainfall over the catchment, SST and output from global climate models.